

The Knaster-Tarski fixed point theorem and the Cantor-Schröder-Bernstein Theorem

Naproche formalization:

Alexander Holz, Marcel Schütz

2024

This is a formalization of the *Knaster-Tarski Fixed Point Theorem* (i.e. of the assertion that every subset-preserving map has a fixed point) and of the *Cantor-Schröder-Bernstein Theorem* (i.e. of the fact that two sets are equinumerous iff they can be embedded into each other), where the proof of the latter theorem is based on the former one, as in [Schroeder2012].

```
[readtex libraries/source/foundations/equinumerosity.ftl.tex]
[readtex libraries/source/foundations/maps-and-systems-of-sets.
ftl.tex]
[readtex libraries/source/set-theory/zf.ftl.tex]
```

The Knaster-Tarski Fixed Point Theorem

Theorem (Knaster-Tarski). Let x be a set. Let h be a map from $\mathcal{P}(x)$ to $\mathcal{P}(x)$ that preserves subsets. Then h has a fixed point.

Proof. (1) Define $A = \{y \mid y \subseteq x \text{ and } y \subseteq h(y)\}$. Then A is a subset of $\mathcal{P}(x)$. We have $\bigcup A \in \mathcal{P}(x)$.

Let us show that (2) $\bigcup A \subseteq h(\bigcup A)$. Let $u \in \bigcup A$. Take $y \in A$ such that $u \in y$. Then $u \in h(y)$. We have $y \subseteq \bigcup A$. Hence $h(y) \subseteq h(\bigcup A)$. Thus $h(y) \subseteq h(\bigcup A)$. Therefore $u \in h(\bigcup A)$. End.

Then $h(\bigcup A) \in A$ (by 1). Indeed $h(\bigcup A) \subseteq x$. (3) Hence $h(\bigcup A) \subseteq \bigcup A$. Indeed every element of $h(\bigcup A)$ is an element of some element of A .

Thus $h(\bigcup A) = \bigcup A$ (by 2, 3). \square

The Cantor-Schröder-Bernstein Theorem

Theorem (Cantor-Schröder-Bernstein). Let x, y be sets. x and y are equinumerous iff there exists a injective map from x to y and there exists an injective map from y to x .

Proof. Case x and y are equinumerous. Take a bijection f between x and y . Then f^{-1} is a bijection between y and x . Hence f is an injective map from x to y and f^{-1} is an injective map from y to x . End.

Case there exists an injective map from x to y and there exists an injective map from y to x . Take an injective map f from x to y . Take an injective map g from y to x . We have $y \setminus f[a] \subseteq y$ for any $a \in \mathcal{P}(x)$.

(1) Define $h(a) = x \setminus g[y \setminus f[a]]$ for $a \in \mathcal{P}(x)$.

h is a map from $\mathcal{P}(x)$ to $\mathcal{P}(x)$. Indeed $h(a)$ is a subset of x for each subset a of x .

Let us show that h preserves subsets. Let u, v be subsets of x . Assume $u \subseteq v$. Then $f[u] \subseteq f[v]$. Hence $y \setminus f[v] \subseteq y \setminus f[u]$. Thus $g[y \setminus f[v]] \subseteq g[y \setminus f[u]]$. Indeed $y \setminus f[v]$ and $y \setminus f[u]$ are subsets of y . Therefore $x \setminus g[y \setminus f[u]] \subseteq x \setminus g[y \setminus f[v]]$. Consequently $h[u] \subseteq h[v]$. End.

Hence we can take a fixed point c of h (by [Knaster-Tarski](#)).

(2) Define $F(u) = f(u)$ for $u \in c$.

We have $c = h(c)$ iff $x \setminus c = g[y \setminus f[c]]$. g^{-1} is a bijection between $\text{range}(g)$ and y . Thus $x \setminus c = g[y \setminus f[c]] \subseteq \text{range}(g)$. Therefore $x \setminus c$ is a subset of $\text{dom}(g^{-1})$.

(3) Define $G(u) = g^{-1}(u)$ for $u \in x \setminus c$.

F is a bijection between c and $\text{range}(F)$. G is a bijection between $x \setminus c$ and $\text{range}(G)$.

Define

$$H(u) = \begin{cases} F(u) & : u \in c \\ G(u) & : u \notin c \end{cases}$$

for $u \in x$.

Let us show that H is a map to y . $\text{dom}(H)$ is a set and every value of H is an object. Hence H is a map.

Let us show that every value of H lies in y . Let v be a value of H . Take $u \in x$ such that $H(u) = v$. If $u \in c$ then $v = H(u) = F(u) = f(u) \in y$. If $u \notin c$ then $v = H(u) = G(u) = g^{-1}(u) \in y$. End. End.

(4) H is surjective onto y . Indeed we can show that every element of y is a value of H . Let $v \in y$.

Case $v \in f[c]$. Take $u \in c$ such that $f(u) = v$. Then $F(u) = v$. End.

Case $v \notin f[c]$. Then $v \in y \setminus f[c]$. Hence $g(v) \in g[y \setminus f[c]]$. Thus $g(v) \in$

$x \setminus h(c)$. We have $g(v) \in x \setminus c$. Therefore we can take $u \in x \setminus c$ such that $G(u) = v$. Then $v = H(u)$. End. End.

(5) H is injective. Indeed we can show that for all $u, v \in x$ if $u \neq v$ then $H(u) \neq H(v)$. Let $u, v \in x$. Assume $u \neq v$.

Case $u, v \in c$. Then $H(u) = F(u)$ and $H(v) = F(v)$. We have $F(u) \neq F(v)$. Hence $H(u) \neq H(v)$. End.

Case $u, v \notin c$. Then $H(u) = G(u)$ and $H(v) = G(v)$. We have $G(u) \neq G(v)$. Hence $H(u) \neq H(v)$. End.

Case $u \in c$ and $v \notin c$. Then $H(u) = F(u)$ and $H(v) = G(v)$. Hence $v \in g[y \setminus F[c]]$. We have $G(v) = g^{-1}(v) \in y \setminus F[c]$. Indeed $v = g(v')$ for some $v' \in y \setminus F[c]$. Thus $G(v) \neq F(u)$. End.

Case $u \notin c$ and $v \in c$. Then $H(u) = G(u)$ and $H(v) = F(v)$. Hence $u \in g[y \setminus F[c]]$. We have $G(u) = g^{-1}(u) \in y \setminus F[c]$. Indeed $u = g(u')$ for some $u' \in y \setminus F[c]$. Thus $G(u) \neq F(v)$. End. End.

Consequently H is a bijection between x and y (by 4, 5). Therefore x and y are equinumerous. End. \square

Copyright

This work is licensed under a [Creative Commons](https://creativecommons.org/licenses/by-nc-sa/4.0/) “Attribution-NonCommercial-ShareAlike 4.0 International” license.

