

## Natural Numbers

SET\_THEORY\_03\_4310076227584000

**Definition 0.1.**

$$\omega = \left\{ n \in \mathbf{Ord} \mid \begin{array}{l} n \in X \text{ for every } X \subseteq \mathbf{Ord} \text{ such that } 0 \in X \text{ and} \\ \text{for all } x \in X \text{ we have } \text{succ}(x) \in X \end{array} \right\}.$$

Let a natural number stand for an element of  $\omega$ .

SET\_THEORY\_03\_3576717620805632

**Proposition 0.2.**  $0 \in \omega$ .

SET\_THEORY\_03\_8807317141192704

**Proposition 0.3.** Let  $n \in \omega$ . Then  $\text{succ}(n) \in \omega$ .

SET\_THEORY\_03\_344585425387520

**Proposition 0.4.** Let  $\Phi \subseteq \omega$ . Assume that  $0 \in \Phi$  and for every  $x \in \Phi$  we have  $\text{succ}(x) \in \Phi$ . Then  $\Phi = \omega$ .

*Proof.* Suppose  $\Phi \neq \omega$ . Consider an element  $n$  of  $\omega$  that is not contained in  $\Phi$ . Take  $\Phi' = \Phi \setminus \{n\}$ .

(1)  $0 \in \Phi'$ . Indeed  $0 \in \Phi$  and  $0 \neq n$ .

(2) For each  $x \in \Phi'$  we have  $\text{succ}(x) \in \Phi'$ .

*Proof.* Let  $x \in \Phi'$ . Then  $\text{succ}(x) \in \Phi$ .

Let us show that  $\text{succ}(x) \neq n$ . Assume  $\text{succ}(x) = n$ . Then  $x \notin \Phi$ . Indeed  $n \notin \Phi$  and if  $x \in \Phi$  then  $n = \text{succ}(x) \in \Phi$ . Contradiction. End.

Thus  $\text{succ}(x) \in \Phi'$ . Qed.

Therefore every element of  $\omega$  lies in  $\Phi'$ . Indeed  $\Phi' \subseteq \mathbf{Ord}$ . Consequently  $n \in \Phi'$ . Contradiction.  $\square$

SET\_THEORY\_03\_4847727433220096

**Corollary 0.5.**  $\omega$  is a set.

*Proof.* Define  $f(n) = \text{succ}(n)$  for  $n \in \omega$ . Take a subset  $X$  of  $\omega$  that is inductive regarding 0 and  $f$ . Indeed  $f$  is a map from  $\omega$  to  $\omega$ . Then we have  $0 \in X$  and for each  $n \in X$  we have  $\text{succ}(n) \in X$ . Thus  $X = \omega$ . Therefore

$\omega$  is a set.

□

SET\_THEORY\_03\_5885789275684864

**Proposition 0.6.** Let  $n \in \omega$ . Then  $n = 0$  or  $n = \text{succ}(m)$  for some  $m \in \omega$ .

*Proof.* Assume the contrary. Consider a  $k \in \omega$  such that neither  $k = 0$  nor  $k = \text{succ}(m)$  for some  $m \in \omega$ . Take a class  $\omega'$  such that  $\omega' = \omega \setminus \{k\}$ . Then  $\omega'$  is a set.

(1)  $0 \in \omega'$ . Indeed  $k \neq 0$ .

(2) For all  $m \in \omega'$  we have  $\text{succ}(m) \in \omega'$ .

Proof. Let  $m \in \omega'$ . Then  $\text{succ}(m) \neq k$ . Hence  $\text{succ}(m) \in \omega'$ . Qed.

Thus every element of  $\omega$  is contained in  $\omega'$ . Therefore  $k \in \omega'$ . Contradiction. □

SET\_THEORY\_03\_5057540872208384

**Proposition 0.7.** Every element of  $\omega$  is an ordinal.

SET\_THEORY\_03\_1991057988386816

**Definition 0.8.**  $1 = \text{succ}(0)$ .

SET\_THEORY\_03\_5809204518453248

**Definition 0.9.**  $2 = \text{succ}(1)$ .

SET\_THEORY\_03\_4388003120152576

**Proposition 0.10.**  $1 = \{0\}$ .

SET\_THEORY\_03\_930896899211264

**Proposition 0.11.**  $2 = \{0, 1\}$ .